

Implementations of Nonadiabatic Geometric Quantum Computation using NMR

Jiangfeng Du,^{1,*} Mingjun Shi,¹ Jihui Wu,² Xianyi Zhou,¹ and Rongdian Han¹

¹*Department of Modern Physics, University of Science and Technology of China, Hefei, 230027, People's Republic of China.*

²*Laboratory of Structure of Biology, University of Science and Technology of China, Hefei, 230027, People's Republic of China.*

Recently, geometric phases, which is fault tolerate to certain errors intrinsically due to its geometric property, are getting considerable attention in quantum computing theoretically. So far, only one experiment about adiabatic geometric gate with NMR through Berry phase has been reported. However, there are two drawbacks in it. First, the adiabatic condition of Berry phase makes such gate very slowly. Second, the extra operation to eliminate the dynamic phase. As we know, geometric phase can exist both adiabatic(Berry phase) and nonadiabatic(Aharonov-Anandan phase). In this letter, we reports the first experimental realization of nonadiabatic geometric gate with NMR through conditional-AA phase. In our experiment the gates can be made faster and more easily, and the two drawbacks mentioned above are removed.

Quantum computers can perform certain tasks much more efficiently than classical Turing Machine[1]. It is well known that controlled two-qubit gate, combined with single qubit operations, is a universal gate for quantum computation[2]. This two-qubit gate preserves the target qubit for the controlling qubit in certain state, say, $|\uparrow\rangle$, and flips the target qubit for the controlling qubit in the other state, $|\downarrow\rangle$. Originally this has been achieved experimentally using dynamic method in different physical systems[3, 4, 5].

On the other hand, central to the experimental realization of quantum computer is the construction of fault-tolerant quantum logic gates[6]. In the quest for a low noise quantum computing device, geometric phases[7, 8], which is fault tolerate to certain errors intrinsically due to its geometric property, are getting considerable attention in quantum computing theoretically[9, 10, 11, 12, 13, 14, 15, 16, 17, 18]. Geometric logical gates based on Berry phase[7] restricted in adiabatic evolution have been proposed[10, 11], and the first adiabatic geometric gate was implemented in NMR[9]. However, the experiments rely on the adiabatic operations. This is a bit impractical because the experimental result is inexact unless the Hamiltonian changes extremely slowly in the process. But in fact, everything has to be completed within the decoherence time. Besides the adiabatic condition, both of the previous proposals require extra operation to eliminate the dynamic phase. This extra operation is unwanted for a fault tolerate gate because if we can not eliminate the dynamic phase exactly, the fault tolerate property is weakened. For these reasons, one is tempted to set up a new scheme which does not rely on the adiabatic condition and which does not involve any accessory operation performed to eliminate dynamic phase in the whole process.

Indeed, geometric phase does exist in a non-adiabatic process. It was shown by Aharonov and Anandan that the geometric phase for a two-level system is only dependent on the area enclosed by the loop on the Bloch sphere[8]. In the non-adiabatic case, the path of the state evolution is, in general, different from the path of the pa-




FIG. 1: The path evolution of qubit a when qubit b is in $|\uparrow\rangle$ state. The first pulse (Fig.2) transformed the initial Hamiltonian $2\pi J I_z^a$ to $H = 2\pi J (I_z^a \cos \theta - I_y^a \sin \theta)$ which creates an evolution path on the geodesic circle ABC. After time $\tau = 1/(2J)$, the second pulse changes the Hamiltonian to the form $H = 2\pi J (-I_z^a \cos \theta - I_y^a \sin \theta)$ which creates an evolution path on the geodesic circle CDA. Again after time $\tau = 1/(2J)$, the third pulse restored the Hamiltonian to the initial form $2\pi J I_z^a$. Therefore, qubit a undergoes a cyclic evolution through a slice circuit C with angle θ in projective Hilbert (density operator) space, and then the AA geometrical phase is simply $\beta(C) = m \Omega$, where $m = \pm \frac{1}{2}$ is the magnetic quantum number and $\Omega = 4\theta$ is the solid angle subtended by the slice circuit.

rameters in the Hamiltonian. The external field need not always follow the evolution path of the state, in contrast to the adiabatic case. Thus, it is possible to let the external field be perpendicular to the evolution path instantaneously so that there is no dynamic phase involved in the whole process. Recently, several schemes were proposed for nonadiabatic geometric gates in different systems[19, 20, 21, 22, 23], but there have been no corresponding experimental demonstrations. Here we show the realization of nonadiabatic two-qubit gate through conditional-AA phase of one two-level subsystem (qubit) controlled by the state of another qubit.

Experimentally, the controlled two-qubit gate was im-




FIG. 2: Pulse sequence used to demonstrate controlled-AA phase of the state of qubit a or controlled two-qubit gate. The black boxes are pulses oscillating at frequency $\omega'_a = \omega_a - \pi J$, the flip angles are $-\theta$, $2\theta - \pi$ and $\pi - \theta$ from left to right which can be realized by choosing the different pulse duration t and pulse power P . Here θ was selected from 0 to π by $\theta = \frac{n\pi}{16}$, $n = \{0, 1, \dots, 16\}$. All pulses that oscillated at frequency $\omega_a - \pi J$ are hard pulses, each pulse duration is $5\mu s$. Delay times are $\tau = 1/2J$ between pair of pulses. Then the time of nonadiabatic controlled two-qubit gate is about $4.8ms$. Note this gate time do not depend on the value of AA phase.

plemented by the nuclear spins of the 1H and ^{13}C atoms in a Carbon-13 labeled chloroform molecule, the single 1H nucleus was used as target qubit a , while the ^{13}C nucleus was used as controlled qubit b , $|\uparrow\rangle(|\downarrow\rangle)$ describes the spin state aligned with (against) an externally applied, strong static magnetic field B_0 in the $\hat{z}$ direction. The reduced Hamiltonian for this two-spin system is, to an excellent approximation, given by $H = \omega_a I_z^a + \omega_b I_z^b + 2\pi J I_z^a I_z^b$, where the first two terms describe the free precession of spin a (1H) and b (^{13}C) about B_0 with frequencies $\omega_a/2\pi \approx 500MHz$ and $\omega_b/2\pi \approx 125MHz$. I_z^a (I_z^b) are the angular momentum operator in the $\hat{z}$ direction for a (b). The third term of the Hamiltonian describes a scalar spin-spin coupling of the two spins with $J = 214.9Hz$. The spin-spin relaxation times are $0.3s$ for carbon and $0.4s$ for proton, respectively. Realistically, the state of this system is a thermally equilibrium one, while pure 00 state must be prepared for most quantum computing tasks. To solve this problem, a variety of techniques exist to extract from this thermal state just the signal from the molecules in the 00 state[25, 26, 27], i.e. all the spins aligned in $+z$ direction. Here we adopt the method of “spatial averaging” to create this effective pure 00 state[25]. The initial state of spin a is $|+\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle)$, the start point of the path evolution, which is prepared by a 90° pulse oscillating at frequency ω_a along y -axes. This resonance pulse rotate the pure state of qubit a from $+z$ -axes to $+x$ -axes.

To produce controlled-AA phase of the state, we change the oscillating frequency ω'_a of qubit a from ω_a to $\omega_a - \pi J$. Now the Hamiltonian of qubit a can be written as $H_a^{rot} = (\omega_a - \omega'_a \pm \pi J) I_z^a = (\pi J \pm \pi J) I_z^a$, which describes the qubit rotating around $+z$ -axis in rotational frame with angular velocity $\omega'_a = \omega_a - \pi J$. That is, when qubit b is in the state $|\uparrow\rangle$, $H_a^{rot} = 2\pi J I_z^a$; when qubit b in $|\downarrow\rangle$, $H_a^{rot} = 0$. Consequently, different states of qubit b

correspond to different frequencies of qubit a in rotational frame. It is just this property that we will use to realize ‘controlled-AA phase shift’. In Fig.1, we show the cyclic evolution of 1H nucleus (qubit a) on the Bloch sphere when the ^{13}C nucleus (qubit b) is in the state $|\uparrow\rangle$. The pulse sequences to realize this nonadiabatic cyclic evolution was shown in Fig.2. As we know, in toggling frame, the track of Hamiltonian is defined by $H = H(0) + H1(t)$, where $H(0)$ is reduced Hamiltonian of the qubit a ($(\pi J \pm \pi J) I_z^a$), and $H1(t)$ is Hamiltonian of RF pulses. Therefore, by applying the pulse with angle θ along x -axes (first black box in Fig.2), the Hamiltonian is changed to $H = 2\pi J (I_z^a \cos \theta - I_y^a \sin \theta)$ in toggling frame. Subject to this Hamiltonian, qubit a undergoes precession from initial state $\frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle)$. After time $\tau = 1/(2J)$, the initial state becomes $|\varphi(\tau)\rangle_a = \frac{e^{-i(\pi/2+\theta)}}{\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle)$. Seen from the resulting state $|\varphi(\tau)\rangle_a$, the direction of spin a is along $-x$ -axis, besides a global phase factor. Now by another pulse with angle $2\theta - \pi$ the Hamiltonian of qubit a has the form $H = 2\pi J (-I_z^a \cos \theta - I_y^a \sin \theta)$. Again, subject to this Hamiltonian, qubit a undergoes precession from state $|\varphi(\tau)\rangle_a$. After time $\tau = 1/(2J)$, the state becomes $|\varphi(2\tau)\rangle_a = \frac{e^{-2i\theta}}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle)$. The direction of spin a returns to $+x$ -axis and a phase -2θ appears. Finally we restore Hamiltonian of qubit a to the initial form $2\pi J I_z^a$ by the last pulse with angle $\pi - \theta$. Note that this pulse has no influence on the final state of qubit a , i.e. $|\varphi(2\tau)\rangle_a$.

From the whole process described above, the Hamiltonian of qubit a has experienced a cyclic evolution because of the cyclic property of the pulse sequence. Correspondingly, an evolution path of ABCDA on the Bloch sphere is produced for qubit a , that is $\frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle) \rightarrow \frac{e^{-i(\pi/2+\theta)}}{\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle) \rightarrow \frac{e^{-2i\theta}}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle)$. Obviously the angle θ is the geometric parameter describing the behavior of the Hamiltonian evolution. This geometric parameter also describes the phase difference between the initial and final state. Note that the dynamic phases appearing in two steps of precession cancel out each other. The geometric AA phase is equal to $\beta(C) = -2\theta$. So far we have considered the case in which the state of qubit b is $|\uparrow\rangle$. If qubit b is in $|\downarrow\rangle$, the Hamiltonian of qubit a is zero and nothing will happen to it. In other words, qubit a will preserve itself in this case. Therefore, the time evolution operator of the pulse sequence (Fig.2) has the property $U(2\tau)|\pm\rangle_a|\uparrow\rangle_b = e^{\pm i\beta(C)}|\pm\rangle_a|\uparrow\rangle_b$ and $U(2\tau)|\pm\rangle_a|\downarrow\rangle_b = |\pm\rangle_a|\downarrow\rangle_b$, where $|\pm\rangle$ corresponds to point A and C, respectively, in the Bloch sphere. Hence we can regard qubit b as a controlling qubit and qubit a as the target qubit; controlled-AA phase of qubit a is produced depended on the state of qubit b . In the basis of $|\uparrow\rangle$ and $|\downarrow\rangle$, the unitary operator that describes this




FIG. 3: Experimental values for the controlled-AA phase $\beta(C)$ as a function of solid angle $\Omega = 4\theta$. Experimental points are shown as small squares; theoretical values are shown as smooth curves. We can see the experiment result fit the theory quite well; the remaining errors could be due to phasing process which is influenced by machine noise and non-ideal baseline. Besides, pulse imperfect and relaxation also have contributions.

circle evolution is

$$\begin{pmatrix} \cos(\beta(C)) & i \sin(\beta(C)) & 0 & 0 \\ i \sin(\beta(C)) & \cos(\beta(C)) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

In particular, when $|\beta(C)| = \pi/2$, this gate is just the C-not gate.

In order to measure the overall AA phase $\beta(C)$ of qubit a we also apply a 90° pulse, oscillating at frequency ω_a along y-axes, to transform qubit b in a coherent superposition of states $|\psi(0)\rangle_b = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle)_b$ before the cyclic evolution. Therefore after this cyclic evolution the final state is $\frac{1}{\sqrt{2}}|+\rangle_a(|\uparrow\rangle + |\downarrow\rangle)_b \rightarrow \frac{1}{\sqrt{2}}|+\rangle_a(|\uparrow\rangle + e^{-i\beta(C)}|\downarrow\rangle)_b$. Here, the unobservable AA phase $\beta(C) = 2\theta = \frac{1}{2}\Omega$ of qubit a transfers to inner phase of qubit b (^{13}C in our experiments) which can be observed with NMR. To do so, we use the signal of initial $\frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle)_b$ state as a reference one, compared with the signal of the final $\frac{1}{\sqrt{2}}(|\uparrow\rangle + e^{-i\beta(C)}|\downarrow\rangle)_b$ state; both of them are in-phase doublet but out of phase by $\beta(C)$. Experimentally, we apply additional phasing factor $\beta(C)$ to obtain absorptive lineshape after Fourier transformation.

All experiments are performed at room temperature and pressure on Bruker Avance DMX-500 spectrometer in Laboratory of Structure Biology, University of Science and Technology of China. The experimental results are shown in Fig.3.

Therefore, we have observed controlled-AA phase or implemented a nonadiabatic two-qubit gate. Note that though the observation of AA phase has been done in NMR with a three-level system[24], it cannot be used to implement universal two-qubit gate.

Our experiment resolves two drawbacks of the adiabatic geometric computation, namely the slow evolution and the need of refocusing to eliminate the dynamical phases. Let us now compare the gate time of this nonadiabatic geometric gate to that of adiabatic geometric gate and dynamic gate. Since the gate time is limited directly by the strength of coupling constant J of the sample. Two experiments we selected to compare used the same sample as ours[5, 9], that is, Carbon-13 labelled chloroform sample. In our experiment, it took about 4.8ms to realize the gate, slightly longer than the time used to realize dynamic two-qubit gate (about 2.4ms)[5], yet much shorter than the time it took to realize the adiabatic geometric two-qubit gate (about 120ms)[9]. As the adiabatic geometric gate operates for a significantly longer time, it is much more severely affected by decoherence. This has serious implications for the physical realization of adiabatic geometric quantum computation. On the other hand, since the state is always perpendicular to the effective magnetic field, there is no dynamical phase accumulation during the evolution, hence the resulted phase factor after cyclic evolution was pure geometric phase. Although this nonadiabatic geometric gate is experimentally realized in the NMR system, the basic idea is general, and could be applied in other physical systems. We believe our experiment has led the idea of geometric quantum computation much more practical than before.

We thank Y.D. Zhang, Z.B. Chen and S.Massar for discussion. This work was supported by NSFC and CAS. Part of the ideas were originated while J.-F. Du was visiting Service de Physique Theorique, Universite Libre de Bruxelles, Bruxelles. S. Massar and N. Cerf are gratefully acknowledged for their invitation and hospitality. J.-F. Du thanks X.-W. Zhu for the loan of the chloroform sample.

* Electronic address: djf@ustc.edu.cn

- [1] D. Deutsch, Proc. R. Soc. London A **400**, 97 (1985).
- [2] Barenco, C. H. Bennett, R. Cleve, D. P. DiVincenzo, N. Margolus, P. Shor, T. Sleator, J. A. Smolin, and H. Weinfurter, Phys. Rev. A **52**, 3457 (1995).
- [3] C. Monroe, D. M. Meekhof, B. E. King, W. M. Itano, and D. J. Wineland, Phys. Rev. Lett. **75**, 4714 (1995).
- [4] Q. A. Turchette, C. J. Hood, W. Lange, H. Mabuchi and H. J. Kimble, Phys. Rev. Lett. **75**, 4710 (1995).
- [5] N. A. Gershenfeld and I. L. Chuang, Science **275**, 350 (1997).
- [6] M. Steane, Nature (London) **399**, 124 (1999).
- [7] M. V. Berry, Proc. R. Soc. A. **392**, 45 (1984).
- [8] Y. Aharonov and J. Anandan, Phys. Rev. Lett. **58**, 1593 (1987).
- [9] J.A.Jones, V.Vedral, A. Ekert, and G. Castagnoli, Nature (London) **403**, 869 (2000).
- [10] G. Falci, R. Fazio, G.M.Palma, G. Slewert and V.Vedral, Nature (London) **407**, 355 (2000).

- [11] L.-M.Duan, J.I.Cirac and P.Zoller, *Science* **292**, 1695 (2001).
- [12] T. Pellizzari, S. A. Gardiner, J. I. Cirac, and P. Zoller, *Phys. Rev. Lett.* **75**, 3788 (1995).
- [13] D. V. Aervin, *Solid State Commun.* **105**, 659 (1998).
- [14] P.Zanardi, *Physical Review A* **63**, 012301 (2001).
- [15] J. Pachos and S. Chountasis, *Physical Review A* **62**, 052318 (2000).
- [16] J. Pachos and M.Rasetti, *Phys. Lett. A* **264**, 94 (1999).
- [17] Pachos, P.Zanardi and M.Rasetti, *Physical Review A* **61**, 010305(R) (2000).
- [18] D. Ellinas and J. Pachos, *Physical Review A* **64**, 022310 (2001).
- [19] W.Xiangbin and M.Keiji, *Physical Review Letter* **87**, 097901 (2001).
- [20] C.Wellard, L.C.L.Hollenberg, and H.C.Pauli, *Physical Review A* **65**, 032303 (2002).
- [21] W.Xiangbin and M.Keiji, *J. Phys. A: Math. Gen.* **34**, L631 (2001).
- [22] W.Xiangbin and M.Keiji, *quant-ph/0104127*.
- [23] Xin-Qi Li et al., *quant-ph/0204028*.
- [24] D. Suter, K. T. Mueller, and A. Pines, *Phys. Rev. Lett.* **60**, 1218 (1988).
- [25] D. G. Cory, M. D. Price, and T. F. Havel, *Physica D* **120**, 82 (1998).
- [26] E. Knill, I. Chuang, and R. Laflamme, *Phys. Rev. A* **57**, 3348 (1997).
- [27] L. M. K. Vandersypen, C. S. Yannoni, M. H. Sherwood, and I. L. Chuang, *Phys. Rev. Lett.* **83**, 3085 (1999).